arXiv:1303.6043v1 [quant-ph] 25 Mar 2013

Local oscillator fluctuation opens a loophole for Eve in practical continuous-variable quantum key distribution system

Xiang-Chun Ma,[1] Shi-Hai Sun,[1] Mu-Sheng Jiang,[1] and Lin-Mei Liang[1, 2, *]

[1]*Department of Physics, National University of Defense Technology, Changsha 410073, P.R.China*
[2]*State Key Laboratory of High Performance Computing,*
National University of Defense Technology, Changsha 410073, P.R.China

We consider the security of practical continuous-variable quantum key distribution implementation with local oscillator (LO) fluctuating in time, which opens a loophole for Eve to intercept the secret key. We show that Eve can simulate this fluctuation to hide her Gaussian collective attack by reducing the intensity of LO. Numerical simulations demonstrate that, if Bob don't monitor LO intensity and scale his measurements with instantaneous intensity values of LO, the secret key rate will be compromised severely.

PACS numbers: 03.67.Dd, 03.67.Hk, 89.70.Cf

I. INTRODUCTION

Continuous-variable quantum key distribution (CVQKD), as an unconditional secure communication scheme between two legitimate parties Alice and Bob, has achieved advanced improvements in theoretical analysis and experimental implementation in recent years [1–5]. Practical implementation systems, such as Fiber-based Gaussian-modulated [6–9] and Discrete-modulated [10, 11] coherent-state protocol QKD systems over tens of kilometers, have been demonstrated in a few groups. The unconditional security of such systems with prepare-and-measure (PM) implementation is confirmed by the security analysis of equivalent entanglement-based (EB) scheme [12–14].

However, the traditional security analysis of EB scheme of CVQKD just includes signal beam, not including local oscillator (LO) which is an auxiliary light beam used as a reference to define the phase of the signal state and is necessary for balanced homodyne detection (BHD). This will leave some security loopholes for Eve because LO is also unfortunate within the Eve's manipulating domain. The necessity of monitoring LO intensity for the security proofs in discrete QKD protocols embedded in continuous variables has been discussed [15]. Moreover, in [16], the excess noise caused by imperfect subtraction of BHD in the presence of LO intensity fluctuations has been noted and quantified with formulation. However, in practical implementation of CVQKD, shot noise scaling with LO power measured before keys distribution is still assumed to keep constant if the fluctuations of LO intensity are small. And in this circumstance pulses with large fluctuation are just discarded as shown in [16]. Unfortunately, this will give Eve some advantages to explore the fluctuation of LO.

In this paper, we primarily analyze the effect of this intensity fluctuation of LO on the secret key rate Alice and Bob hold. To begin the analysis, we first describe Bob's measurements under the fluctuation of LO intensity, and propose an attacking scheme exploring this fluctuation. We consider the security of practical CVQKD implementation under this attack and calculate the secret key rate with and without Bob monitoring LO for reverse and direct reconciliation protocol. We find that the fluctuation of LO could compromise the secret keys severely if Bob don't scale his measurements with instantaneous intensity values of LO. Finally, we briefly discuss the accurate monitoring of LO intensity to confirm the security of practical implementation of CVQKD.

II. LOCAL OSCILLATOR INTENSITY FLUCTUATION AND ATTACK

Generally, in practical system of CVQKD, the local oscillator intensity is always monitored by splitting a small part with a beam splitter and pulses with large intensity fluctuation of LO are discarded too. However, even monitoring, we aren't clear about how the fluctuation, in particular the small fluctuation, affects the secret key rate yet. In order to confirm the secret key rate obtained by Alice and Bob being unconditional secure, in what follows, we will analyze the effects of this fluctuation on secret key rate only, and don't consider the imperfect measurement of BHD due to incomplete subtraction of it in presence of LO intensity fluctuations, which has been discussed in [16].

Ideally, with strong LO, a perfect BHD measuring a weak signal whose quadratures are $X_S \in \{Q_S, P_S\}$ will output the results[17],

$$x_\theta = k|\alpha_{LO}|(Q_{in}\cos\theta + P_{in}\sin\theta), \tag{1}$$

where k is a proportional constant of BHD, α_{LO} is amplitude of LO, θ is the relative phase between signal and LO except signal's initial modulation phase. So scaling with LO power or shot noise, the results can be recast as

$$\hat{X}_O = \hat{X}_{in} = X_S + \hat{X}_N, \tag{2}$$

---

[*] Electronic address: nmliang@nudt.edu.cn

with θ in Eq. (1) is 0 or $\pi/2$. Here the quadratures Q and P are defined as $\hat{X}_{in} \in \{\hat{Q}_{in}, \hat{P}_{in}\}$ and $\hat{X}_N \in \{\hat{Q}_N, \hat{P}_N\}$, where $\hat{X}_N$ is the quadrature of vacuum state.

However, in practical system, the intensity of LO fluctuates in time. With a proportional coefficient $\eta > 0$, practical LO intensity can be described as $|\alpha'_{LO}|^2 = \eta|\alpha_{LO}|^2$, where α_{LO} is the initial amplitude of LO used by normalization. If we don't monitor LO or quantify its fluctuation, especially just let the outputs of BHD scale with the initial intensity or power of LO, the outputs then read

$$\hat{X}_O^{'} = \sqrt{\eta}\hat{X}_O. \tag{3}$$

Unfortunately, this fluctuation will open a loophole for Eve, as we will see in the following sections.

In conventional security analysis, like EB scheme equivalent to usual PM implementation depicted in Fig. 1(a), LO is not taken into consideration and its intensity is assumed to keep unchanged. However, in practical implementation, Eve not only intercept the signal beam but also LO and she can replace the quantum channel between Alice and Bob with her own perfect quantum channel as shown in Fig. 1(b) and (c). In so doing, Eve's attack can be partially hidden by reducing the intensity of LO with an attenuator simulating the fluctuation without changing LO's phase, and such attack can be called LO intensity attack (LOIA). In following analysis, we will see that, in parameter estimation between Alice and Bob, channel excess noise introduced by Eve can be reduced arbitrarily, even to be null, with Eve just by tuning the transmission of LO. Consequently, Alice and Bob would underestimate Eve's intercepted information and Eve could get partial secret keys Alice and Bob hold without being found under this attack.

Fig. 1(b) describes the LOIA, which consists in attacking the signal beam with general Gaussian collective attack[12–14] and attacking LO beam with intensity attenuation by a no changing phase attenuator **A**, such as a beam splitter whose transmission is variable. This signal beam Gaussian collective attack consists of three steps: Eve interacts her ancilla modes with the signal mode by a unitary operation **U** for each pulse and store them in her quantum memory, then she make an optimal collective measurement after Alice and Bob's classical communication. Fig. 1(c) is one practical LOIA with **U** being a beam splitter transformation. Its signal attack is also called entangling cloner attack, which is presented first by Grosshan [18] and improved by Weedbrook [19, 20]. We will give a demonstration that with this entangling cloner Eve can get the same amount of information as much as the one Fig. 1(b) shows in appendix A.

III. ESTIMATION OF SECRET KEY RATE UNDER LOIA

We analyze a practical CVQKD system with homodyne protocol for example to demonstrate the effect of



FIG. 1: (a) EB scheme, based on Alice heterodyning one of the mode of Einstein-Podolsky-Rosen(EPR)[23] state and Bob homodyning the other half, with Eve only attacking the signal. (b) LOIA consisting in attacking the signal beam with general Gaussian collective attack and attacking LO with intensity attenuation by a no changing phase attenuator **A** such as a beam splitter whose transmission is variable. **U** is unitary operation and X_E is Eve's intercepted mode. (c) LOIA with **U** being a beam splitter transformation and Eve's ancilla modes X_{E0} and X_{E2} being the EPR pairs with variance N, X_{E1} intercepted mode. This signal attack is also called entangling cloner attack.

LOIA on secret key rate, and for simplicity we don't give the results of heterodyne protocol which is analogous with homodyne protocol. In usual PM implementation, Alice prepares a series of coherent states centered on $X_S \in \{Q_S, P_S\}$ with each pulse, Q_S, P_S respectively satisfy a Gaussian distribution independently with the same variance V_S and zero mean, and she then send them to Bob through a quantum channel which might be intercepted by Eve. This initial mode prepared by Alice can be described as $\hat{X}_A = X_S + \hat{X}_N$, and $\hat{X}_A \in \{\hat{Q}_A, \hat{P}_A\}$ is the quadrature variable. Here $\hat{X}_N \in \{\hat{Q}_N, \hat{P}_N\}$ describes the quadratures of vacuum mode. Note that we denote an operator with a hat, while without a hat the same variable corresponds to the classical variable after measurement. So the overall variance of initial mode prepared by Alice is $V = V_S + 1$. When this mode comes to Bob, Bob will get a mode $\hat{X}_B \in \{\hat{Q}_B, \hat{P}_B\}$,

$$\hat{X}_B = \sqrt{T}\hat{X}_A + \sqrt{1-T}\hat{E}, \tag{4}$$

where $\hat{E}$ describes Eve's mode introduced through the quantum channel whose quadrature variance is N. Bob randomly selects a quadrature to measure, and if Eve attenuates the intensity of LO, but Bob's outputs still scale with the initial LO intensity, just like Eq. (3) he will get the measurement

$$X_B^m = \sqrt{\eta}X_B = \sqrt{\eta}(\sqrt{T}X_A + \sqrt{1-T}E), \tag{5}$$

where X_B is $Q_B \in \mathbb{R}$ (or equivalently$P_B \in \mathbb{R}$). However, if Bob monitors LO and also scales with the instantaneous intensity value of LO with each pulse, he will get

X_B without any loss of course. Note that in this paper we just talk about an optimal measure and the efficiency and other imperfections of Bob's BHD are not considered. Thus the variance of Bob's measurements and conditional variance on Alice's encodings with and without monitoring (in what follows, without monitoring specially indicates that Bob's measurement is obtained just by scaling with initial LO intensity instead of monitoring instantaneous value, and vice versa) can be given, respectively, by

$$\begin{aligned} V_B &= TV + (1-T)N\ , && (6)\\ V_B^m &= \eta\left[TV + (1-T)N\right]\ , && (7)\\ V_{B|A} &= T + (1-T)N\ , && (8)\\ V_{B|A}^m &= \eta\left[T + (1-T)N\right]\ , && (9) \end{aligned}$$

where all variances are in shot noise units and the conditional variance is defined as[21, 22]

$$V_{X|Y} = V(X) - \frac{|\langle XY\rangle|^2}{V(Y)}. \qquad (10)$$

Hence, the covariance matrix of Alice's and Bob's mode can be obtained as

$$\begin{aligned} \gamma_{AB}(V,T,N) &= \begin{pmatrix} \gamma_A & \sigma_{AB}^T \\ \sigma_{AB} & \gamma_B \end{pmatrix} \\ &= \begin{pmatrix} V\mathbb{I} & \sqrt{T(V^2-1)}\sigma_z \\ \sqrt{T(V^2-1)}\sigma_z & (TV+(1-T)N)\cdot\mathbb{I} \end{pmatrix}, \qquad (11) \\ \gamma_{AB}^m &= \begin{pmatrix} V\mathbb{I} & \sqrt{\eta T(V^2-1)}\sigma_z \\ \sqrt{\eta T(V^2-1)}\sigma_z & \eta\left[TV+(1-T)N\right]\mathbb{I} \end{pmatrix}, \qquad (12) \end{aligned}$$

where $\sigma_z = \left(\begin{smallmatrix} 1 & 0 \\ 0 & -1 \end{smallmatrix}\right)$ is Pauli matrix and $\mathbb{I}$ is unit matrix.

From Eq. (6) and (7) we can derive that the channel transmission and excess noise are T, $\varepsilon = (1-T)(N-1)/T$ with monitoring, and ηT, $\varepsilon^m = \varepsilon - \frac{1}{T}(\frac{1}{\eta}-1)$ without monitoring, respectively. Hence, by attenuating LO intensity as Fig. 1 shows to make $0 < \eta < 1$, Eve could arbitrarily reduce ε^m to zero, thus she will get the largest amount of information permitted by physics. In following numerical simulation we always make $\varepsilon^m = 0$, namely, $\eta(1-T)N = 1 - \eta T$. Thus, the covariance matrix $\gamma_{AB}^m = \gamma_{AB}(V, \eta T, 1)$ and Eve's introducing noise N (in entangling cloner it is Eve's EPR state's variance as Fig. 1(c) shows) should be selected to be

$$N = \frac{1-\eta T}{\eta(1-T)}. \qquad (13)$$

In order to estimate the secret key rate, without loss of generality, we first analyze the reverse reconciliation then consider the direct reconciliation.

A. Reverse reconciliation

From Alice and Bob's points of view, the secret key rate for reverse reconciliation with monitoring or not are given, respectively, by

$$\begin{aligned} K_{RR} &= I_{AB} - \chi_{BE}, && (14)\\ K_{RR}^m &= I_{AB}^m - \chi_{BE}^m, && (15) \end{aligned}$$

where the mutual information between Alice and Bob with and without monitoring are the same, and that's

$$I_{AB} = \frac{1}{2}\log\frac{V_B}{V_{B|A}} = \frac{1}{2}\log\frac{V_B^m}{V_{B|A}^m} = I_{AB}^m. \qquad (16)$$

This is because Bob's measurements in these two cases are just different with a coefficient η, and they correspond with each other one by one, so they are equivalent according to data processing theorem[24, 25]. However, the mutual information between Eve and Bob given by Holevo bound[26] in these two cases are not identical. As the previous analysis showed, in Bob's point of view, channel transmission and the excess noise estimation are different. But from Eve's point of view, they are identical according to data processing theorem because she estimates Bob's measurements in these two cases just multiplying a coefficient η. We'll calculate the practical information intercepted by Eve first. It can be given by

$$\chi_{BE} = S(E) - S(E|B), \qquad (17)$$

where $S(\cdot)$ is the Von Neumann entropy[27]. For a Gaussian state ϱ, this entropy can be calculated by the symplectic eigenvalues of the covariance matrix γ characterizing ϱ. To calculate Eve's information, first, Eve's system E can purify AB permitted by quantum physics, so that $S(E) = S(AB)$. Second, after Bob's projective measurement, the system AE is pure, so that $S(E|B) = S(A|B)$. Designating $a = V, b = TV + (1-T)N, c = \sqrt{T(V^2-1)}$, the symplectic eigenvalues of γ_{AB} are given by

$$\lambda_{1,2} = \sqrt{\frac{A \mp \sqrt{A^2 - 4B^2}}{2}}, \qquad (18)$$

wher e $A = a^2 + b^2 - 2c^2$ and $B = ab - c^2$. Similarly, the entropy $S(A|B)$ is determined by the symplectic eigenvalue λ_3 of covariance matrix $\gamma_A^{\hat{x}_B}$[29], namely,

$$\gamma_A^{\hat{x}_B} = \gamma_A - \sigma_{AB}^T(\mathbf{X}\gamma_B\mathbf{X})^{MP}\sigma_{AB}, \qquad (19)$$

where $\mathbf{X} = \left(\begin{smallmatrix} 1 & 0 \\ 0 & 0 \end{smallmatrix}\right)$ and MP stands for the Moore Penrose inverse of a matrix. Then $\lambda_3 = \sqrt{\frac{(1-T)NV^2+TV}{TV+(1-T)N}}$, the Holevo bound reads

$$\chi_{BE}(V,T,N) = G\big(\frac{\lambda_1-1}{2}\big) + G\big(\frac{\lambda_2-1}{2}\big) - G\big(\frac{\lambda_3-1}{2}\big), \qquad (20)$$

where $G(x) = (x+1)\log_2(x+1) - x\log_2 x$. However, Eve's information estimated by Bob without monitoring is given by

$$\chi_{BE}^m = \chi_{BE}(V, \eta T, 1). \qquad (21)$$

Substituting Eq. (20) and (21) into Eq. (14) and (15) respectively, the secret key rate with and without Bob's monitoring can be obtained. However, the secret key rate in Eq. (15) without monitoring is unsecure in evidence. Eve intercepted partial information from K^m_{RR} but not be found, in other words, Alice and Bob underestimate Eve's information but without realizing it. Actually, the practical or unconditional secure secret key rate K^m_{RR}, which we called truly secret key rate, should be available by replacing χ^m_{BE} in Eq. (15) with Eq. (20). Note that it is identical with the monitoring secret key rate in Eq. (14) due to Eq. (16).

We investigate the secret key rate K^m_{RR} Bob measured without monitoring and the truly one or equivalently monitoring one K_{RR} for reverse reconciliation under Eve attacking the intensity of LO. As Fig. 2 shows, with various values of transmission of LO that can be controlled by Eve, the truly secret key rate Alice and Bob actually share decreases rapidly over long distances or small channel transmission.



FIG. 2: (Color online)Reverse reconciliation pseudo-secret key rate and truly one vs. channel transmission T under LOIA. Solid lines are secret key rate estimated by Bob without monitoring LO intensity and dashed lines are the truly ones. Color 'red', 'green', 'blue', 'cyan', 'magenta' corresponds to the LO transmission $\eta = 0.8, 0.85, 0.9, 0.95, 1$, respectively. Here Alice's modulation variance $V_S = 20$.

Additionally, because the mutual information between Alice and Bob with and without monitoring are identical as Eq.(16) shows, subtracting Eq. (14) from Eq. (15) we can estimate Eve's intercepted information $(K^m_{RR} - K_{RR})$ which is plotted in Fig. 3. We find that Eve could get partial or full secret keys which Alice and Bob hold by controlling the different transmission of LO. Taking 20km transmission distance for example, surprisingly, just with LO intensity fluctuation or attenuating rate 0.08, Eve is able to obtain the full secret keys for reverse reconciliation without Bob's monitoring LO.



FIG. 3: (Color online)Reverse reconciliation pseudo-secret key rate (dashed lines) and intercepted one (solid lines) by Eve without being found vs. LO intensity fluctuation rate $(1 - \eta)$ with different transmission distances. From top to bottom the transmission distances are 10km, 20km, 30km, 40km, and 50km, respectively. Inset is amplified one with $(1 - \eta)$ between $0 \sim 0.16$. Here Alice's modulation variance $V_S = 20$.

B. Direct reconciliation

We now calculate the secret key rate for direct reconciliation, which is a little more complicated, and investigate the effect of LO intensity attack by Eve on CVQKD. The secret key rate estimated by Bob with and without LO monitoring is given, respectively, by

$$K_{DR} = I_{AB} - \chi_{AE}, \tag{22}$$

$$K^m_{DR} = I^m_{AB} - \chi^m_{AE}. \tag{23}$$

Note that we have already calculated I_{AB} and I^m_{AB} in Eq.(16) and they are identical for direct and reverse reconciliation. For Eve we have

$$\chi_{AE} = S(E) - S(E|A), \tag{24}$$

where $S(E) = S(AB)$ has been already computed in previous section, and $S(E|A) = S(BC|A)$ using the fact that after Alice's projective measurement on mode A_0 and C_0 obtaining Q_A in EB scheme shown in Fig. 1(a), the system BCE is pure. To calculate $S(BC|A)$, we have to compute the symplectic eigenvalues of covariance matrix $\gamma^{\hat{x}_A}_{BC}$, which is obtained by

$$\gamma^{\hat{x}_A}_{BC} = \gamma_{BC} - \sigma^T_{BCA}(\mathbf{X}\gamma_A\mathbf{X})^{MP}\sigma_{BCA}, \tag{25}$$

where γ_{BC} and σ_{BCA} can be read in the decomposition of the matrix

$$\gamma_{BCA} = \begin{pmatrix} \gamma_{BC} & \sigma^T_{BCA} \\ \sigma_{BCA} & \gamma_A \end{pmatrix}, \tag{26}$$

which is available by elementary transformation of matrix (see Fig. 1(a))[6, 29]

$$\gamma_{ACB} = \big(S^{BS}_{A_0C_0} \oplus \mathbb{I}_B\big)^T \gamma_{A_0C_0B} \big(S^{BS}_{A_0C_0} \oplus \mathbb{I}_B\big), \qquad (27)$$

where $\mathbb{I}_B$ is unit matrix. It is obtained by applying homodyning over mode A after mixing A_0 and C_0 with a balanced beam splitter transformation($S^{BS}_{A_0C_0}$). The matrix $\gamma_{A_0C_0B} = \gamma_{A_0B} \oplus \gamma_{C_0}$ and γ_{A_0B} actually is γ_{AB} in Eq. (11), γ_{C_0} is unit matrix. So we can get

$$\gamma_{BC}^{\hat{x}_A} = \begin{pmatrix} b - c^2/(a+1) & 0 & \sqrt{2}c/(a+1) & 0 \\ 0 & b & 0 & -c/\sqrt{2} \\ \sqrt{2}c/(a+1) & 0 & 2a/(a+1) & 0 \\ 0 & -c/\sqrt{2} & 0 & (a+1)/2 \end{pmatrix} \qquad (28)$$

and the symplectic eigenvalues of it

$$\lambda_{4,5} = \sqrt{\frac{C \mp \sqrt{C^2 - 4D}}{2}}, \qquad (29)$$

where $C = \frac{a+bB+A}{a+1}$ and $D = \frac{B(b+B)}{a+1}$. The Holevo bound then reads

$$\chi_{AE}(V,T,N) = \sum_{i=1}^{2} G\big(\frac{\lambda_i - 1}{2}\big) - \sum_{j=4}^{5} G\big(\frac{\lambda_j - 1}{2}\big), \qquad (30)$$

$$\chi_{AE}^m = \chi_{AE}(V, \eta T, 1). \qquad (31)$$

Substituting Eq. (30) and (31) into Eq. (22) and (23) respectively, the secret key rates in these two cases are obtained. We plotted them for channel transmission T with various values of η and find that the difference between pseudo-secret key rate with Bob not monitoring LO and the truly one is still increasing with the channel transmission T becoming smaller.

For Eve, when Bob don't monitor LO, she will get the partial or total secret key rate $(K_{DR}^m - K_{DR})$ without being found by subtracting Eq. (22) from Eq. (23) when she reduces the intensity of LO. In Fig. 5, we plotted pseudo-secret key rate for direct reconciliation and the mutual information overestimated by Alice and Bob. We find that for short distance communication (less 15km or 3dB limit), a small fluctuation of LO intensity could still hide Eve's attack partially or totally.



FIG. 4: (Color online)Direct reconciliation pseudo-secret key rate and truly one vs. channel transmission T under LOIA. Solid lines are secret key rate estimated by Bob without monitoring LO intensity and dashed lines are the truly ones. Color 'red', 'green', 'blue', 'cyan', 'magenta' corresponds to the LO transmission $\eta = 0.6, 0.7, 0.8, 0.9, 1$, respectively. Here Alice's modulation variance $V_S = 20$.



FIG. 5: (Color online)Direct reconciliation pseudo-secret key rate (dashed lines) and intercepted one (solid lines) by Eve without being found vs. LO intensity fluctuation rate $(1-\eta)$ with different transmission distances. From top to bottom the transmission distances are 6km, 8km, 10km, 12km, and 14km, respectively. Here Alice's modulation variance $V_S = 20$.

IV. DISCUSSION AND CONCLUSION

With the analysis from the previous section, we show that reverse reconciliation is more sensible than direct reconciliation about the fluctuation of LO intensity and even with a small fluctuation or attenuation of LO intensity, Eve can get full secret keys, but not being found, that Alice and Bob hold. This is consistent with that channel excess noise has more severe impact on reverse reconciliation than direct reconciliation. Remarkably, LO intensity fluctuation opens a loophole for Eve, especially in case of communication with low channel transmission or over long distance. Consequently, in practical implementation of CVQKD, we must monitor the LO fluctuation carefully and in particular scale the measurements with instantaneous intensity values of LO. However, for reverse reconciliation communication over long distance, very small fluctuation of LO might compromise the secret key rate completely, which presents a big challenge for accurately monitoring LO intensity.

Finally, we point out that in this paper we don't consider the imperfection of BHD such as detection efficiency and incomplete subtraction etc, which may make LO intensity fluctuation have more severe impact on estimation of secret key rate for Alice and Bob.

In conclusion, we have analyzed the effect of LO intensity fluctuation on secret key rate estimation of Alice and Bob for reverse and direct reconciliation. Incredibly, Bob's estimation of secret key rate will be compromised severely without monitoring LO or his measurements not scaling with LO instantaneous intensity values even with monitoring but just discard large fluctuation pulse like in[16]. Furthermore, we have shown that Eve could hide her attack partially by reducing the intensity of LO and even could steal the total secret keys Alice and Bob share without being found by a small attenuation of LO intensity, especially for reverse reconciliation. Finally, we have also briefly discussed the monitoring of LO and pointed out that it would be a challenge for high accurately monitoring.

Acknowledgments

This work is supported by the National Natural Science Foundation of China, Grant No. 61072071. L-M. L. is supported by the Program for New Century Excellent Talents.

Appendix A: Security estimation by Entangling Cloner under LOIA

In this appendix, we calculate the Holevo bound obtained by Eve for direct and reverse reconciliation using Weedbrook's entangling cloner model[19, 20], and then give the secret key rate shared by Alice and Bob under LOIA. We begin the analysis by calculating the Von Neumann entropy of Eve's intercepting state first.

As Fig. 1(c) shows, the entangling cloner consists in Eve replacing the Gaussian quantum channel between Alice and Bob with a beam splitter of transmission T and an EPR pair of variance N. Half of the EPR pair mode E_0 is mixed with Alice's mode in the beam splitter and is sent to Bob to match the noise of the real channel by tuning N. The other half mode E_2 is kept by Eve to reduce the uncertainty on one output of the beam splitter, the mode E_1 which can be read as

$$\hat{X}_{E_1} = -\sqrt{1-T}\hat{X}_A + \sqrt{T}\hat{X}_{E_0}, \tag{A1}$$

where $\hat{X}_{E_0}$ is the quadrature of mode E_0. Thus, the variance of mode E_1 is given by

$$V_{E_1} = (1-T)V + TN, \tag{A2}$$

and the conditional variance $V_{E_1|A}$ can be calculated as, using Eq. (10),

$$V_{E_1|A} = (1-T) + TN. \tag{A3}$$

Hence, Eve's covariance matrix can be obtained as

$$\gamma_E(V,V) = \begin{pmatrix} \gamma_{E_1} & \sigma_{E_1E_2}^T \\ \sigma_{E_1E_2} & \gamma_{E_2} \end{pmatrix} = \begin{pmatrix} diag(V_{E_1}, V_{E_1}) & Z_{E_1E_2}\sigma_z \\ Z_{E_1E_2}\sigma_z & N\mathbb{I} \end{pmatrix}, \tag{A4}$$

where $Z_{E_1E_2} = \sqrt{T(N^2-1)}$ and the notation $diag(,)$ stands for a matrix with the arguments on the diagonal elements and zeros everywhere else. The symplectic eigenvalues of this covariance matrix are given by

$$\lambda_{1,2} = \sqrt{\frac{\Delta \mp \sqrt{\Delta^2 - 4D}}{2}}, \tag{A5}$$

where $\Delta = V_{E_1}^2 + N^2 - 2Z_{E_1E_2}^2, D = (V_{E_1}N - Z_{E_1E_2}^2)^2$. Hence, the Von Neumann entropy of Eve's state is given by

$$S(E) = G(\frac{\lambda_1 - 1}{2}) + G(\frac{\lambda_2 - 1}{2}), \tag{A6}$$

1. Direct reconciliation

For direct reconciliation protocol of CVQKD, the Holevo bound between Eve and Alice is given by Eq. (24), where $S(E)$ has been calculated by Eq. (A6). $S(E|A)$ can be obtained by the conditional covariance matrix

$$\gamma_E^{X_A} = \gamma_E(V=1, V), \tag{A7}$$

and its symplectic eigenvalues are given by

$$\lambda_{3,4} = \sqrt{\frac{A \mp \sqrt{A^2 - 4B}}{2}}, \tag{A8}$$

wher e $A = V_{E_1|A}V_{E_1} + N^2 - 2Z_{E_1E_2}^2, B = (V_{E_1|A}N - Z_{E_1E_2}^2)(V_{E_1}N - Z_{E_1E_2}^2)$. Thus, the conditional entropy is

$$S(E|A) = G(\frac{\lambda_3 - 1}{2}) + G(\frac{\lambda_4 - 1}{2}). \tag{A9}$$

Substituting Eq. (A6) and (A9) into Eq. (24), we can get the mutual information between Alice and Eve,

$$\chi_{AE}(V,T,N) = S(E) - S(E|A). \tag{A10}$$

Under LOIA, Bob's estimation of the Holevo bound without monitoring LO intensity then reads, using Eq. (A10)

$$\chi_{AE}^m = \chi_{AE}(V, \eta T, 1). \tag{A11}$$

With Eq. (A10) and (A11), the secret key rate in Eq. (22) and (23) then can be calculated respectively, and numerically calculation demonstrates that they are consistent with the Fig. 4 perfectly.

2. Reverse reconciliation

The calculation of Holevo bound between Eve and Bob for reverse reconciliation is a bit more complicated. Using Eq. (17), we only need to calculate the conditional entropy $S(E|B)$, which is determined by the symplectic eigenvalues $\lambda_{4,5}$ of covariance matrix $\gamma_E^{\hat{X}_B}$,

$$\gamma_E^{\hat{X}_B} = \gamma_E - \sigma_{E_1E_2B}^T (\mathbf{X}\gamma_B\mathbf{X})^{MP} \sigma_{E_1E_2B}, \tag{A12}$$

wher e $\sigma_{E_1E_2B} = \left(\left\langle \hat{X}_{E_1}\hat{X}_B \right\rangle \mathbb{I}, \left\langle \hat{X}_{E_2}\hat{X}_B \right\rangle \sigma_z\right) = (Z_{E_1B}\mathbb{I}, Z_{E_2B}\sigma_z)$ and $Z_{E_1B} = \sqrt{T(1-T)}(N - V), Z_{E_2B} = \sqrt{1-T}\sqrt{N^2-1}$. Then, $\gamma_E^{\hat{X}_B}$ can be recast as

$$\gamma_E^{\hat{X}_B} = \begin{pmatrix} F & H^T \\ H & G \end{pmatrix} \tag{A12$'$}$$

where $F{=}diag(V_{E_1}{-}\frac{Z^2_{E_1B}}{V_B}, V_{E_1})$, $G{=}diag(N{-}\frac{Z^2_{E_2B}}{V_B}, N)$, and $H{=}diag(Z_{E_1E_2} - \frac{Z_{E_1B}Z_{E_2B}}{V_B}, -Z_{E_1E_2})$. Hence, its symplectic eigenvalues are given by

$$\lambda_{5,6} = \sqrt{\frac{C \mp \sqrt{C2 - 4D'}}{2}}, \tag{A13}$$

where $C = det(F) + det(G) + 2det(H), D' = det(\gamma_E^{\hat{X}_B})$ and $det(\cdot)$ is the determinant of a matrix. So, we get the conditional entropy

$$S(E|B) = G(\frac{\lambda_5 - 1}{2}) + G(\frac{\lambda_6 - 1}{2}), \tag{A14}$$

and then the Holevo bound

$$\chi_{BE}(V, T, N) = S(E) - S(E|B). \tag{A15}$$

Consequently, without monitoring LO intensity, Alice and Bob will give Eve the Holevo bound

$$\chi^m_{BE} = \chi_{BE}(V, \eta T, 1). \tag{A16}$$

Substituting Eq. (A15) and (A16) into Eq. (14) and (15) respectively, the secret key rate with and without Bob's monitoring can be obtained, and for channel transmission T with various values of η, they are numerically demonstrated to be consistent with Fig. 2 perfectly, too.

Hence, it again indirectly confirm that either for direct or reverse reconciliation, the entangling cloner could reach the Holevo bound against the optimal Gaussian collective attack.

[1] C. Weedbrook, S. Pirandola, R. García-Patrón, N. J. Cerf, T. C. Ralph, J. H. Shapiro, and S. Lloyd, Rev. Mod. Phys. **84**, 621 (2012).

[2] F. Grosshans, G. V. Asschee, J. Wenger, R. Brouri, N. Cerf, and P. Grangier, Nature (London) **421**, 238 (2003).

[3] A. Leverrier, E. Karpov, P. Grangier, and N. J. Cerf, New J. Phys. **11**, 115009 (2009).

[4] A. Leverrier and P. Grangier, Phys. Rev. Lett. **102**, 180504 (2009).

[5] Q. D. Xuan, Z. Zhang, and P. L. Voss, Opt. Exp. **17**, 24244 (2009).

[6] J. Lodewyck *et al.*, Phys. Rev. A **76**, 042305 (2007).

[7] B. Qi, L. L. Huang, L. Qian, and H. K. Lo, Phys. Rev. A **76**, 052323 (2007).

[8] S. Fossier, E. Diamanti, T. Debuisschert, A. Villing, R. Tualle-Brouri, and P. Grangier, New J. Phys. **11**, 045023 (2009).

[9] P. Jouguet *et al.*, Opt. Exp. **20**, 14031 (2012).

[10] Z. Zhang and P. L. Voss, Opt. Exp. **17**, 12090 (2009).

[11] Y. Shen, H-X. Zou, L. Tian, P-X Chen, and J-M Yuan, Phys. Rev. A **82**, 022317 (2010).

[12] F. Grosshans, Phys. Rev. Lett. **94**, 020504 (2005).

[13] R. García-Patrón and N. J. Cerf, Phys. Rev. Lett. **97**, 190503 (2006).

[14] M. Navascués, F. Grosshans, and A. Acín, Phys. Rev. Lett. **97**, 190502 (2006).

[15] H. Haseler, T. Moroder, and N. Lütkenhaus, Phys. Rev. A **77**, 032303 (2008).

[16] Y-M. Chi *et al.*, New J. Phys. **13**, 013003 (2010).

[17] M. G. Raymer, J. Cooper, H. J. Carmichael, M. Beck, and D. T. Smithey, J. Opt. Soc. Am. B **12**, 1801 (1995).

[18] F. Grosshans, N. J. Cerf, J. Wenger, R. Tualle-Brouri, and P. Grangier, Quantum Inf. Comput. **3**, 535 (2003).

[19] C. Weedbrook, S. Pirandola, S. Lloyd, and T. C. Ralph, Phys.Rev. Lett. **105**, 110501 (2010).

[20] C. Weedbrook, S. Pirandola, and T. C. Ralph, Phys. Rev. A **86**, 022318 (2012).

[21] J.P. Poizat, J. F. Roch, and P. Grangier, Ann. Phys. (Paris) **19**, 265 (1994).

[22] P. Grangier, J. A. Levenson, and J. P. Poizat, Nature (London) **396**, 537 (1998).

[23] A. Einstein, B. Podolsky, and N. Rosen, Phys. Rev. **47**, 777(1935).

[24] S. Kullback and R. Leibler, Ann. Math. Statist. **22**, 79 (1951).

[25] J. Ziv and M. Zakai, IEEE Trans. Inf. Theory **19**, 275 (1973).

[26] A. S. Holevo, M. Sohma, and O. Hirota, Phys. Rev. A **59**, 1820 (1999).

[27] J. von Neumann, *Mathematical Foundation of Quantum Mechanics* (English Translation by R. T. Beyer, Princeto University Press, Princeton, NJ, 1955).

[28] A. Serafini, F. Illuminati, and S. De Siena, J. Phys. B **37**, L21 (2004); G. Adesso, A. Serafini, and F. Illuminati, Phys. Rev. A **70**, 022318 (2004).

[29] R. García-Patrón, Ph. D. thesis, Université Libre de Bruxelles, 2007.